

# Powerful Performance. Winning Results.



## PRELUDE™309

Honor® Show Chow® PRELUDE $^{\text{M}}$  309 feed is designed to be fed to pigs weighing 16 to 25 lbs, but can be fed to heavier pigs.

Offer PRELUDE™ 309 feed as a phase 2 pig starter when pigs are about 28 days of age or weigh 16 lbs.

Honor® Show Chow® PRELUDE $^{\text{m}}$  309 feed is supported by Purina. Get your newly weaned pigs on PRELUDE $^{\text{m}}$  309 feed today so you can unleash your genetic investment.

FEATURES	BENEFITS	
Formulated with very high quality animal protein sources	Very palatable, newly weaned pigs get on feed fast and look great	
High levels of protein, essential amino acids (1.5% lysine), and crude fat (8.0%)	Nutrient dense to support intake and growth of young pigs	
Organic acids, yucca schdigera extract, and high levels of vitamins.	Acidified to enhance digestibility, yucca to promote intake and nursery air quality, and high in fat and water soluble vitamins for small pigs	
3/32" pellets or crumbles	Extremely small pellet for improved intake and pellet quality	
CTC & Denagard medication	Prevent and control respiratory and digestive tract diseases	
Proprietary blend of ingredients	Produces full middles, muscle shape and bloom	
High lactose levels	Easy transition from sow's milk to solid feed	



Regional Availability

Check your current price list for availability in your region, or contact your local Purina Dealer or Representative for more information.

# PRELUDE™309 CTC/DEN

## **GUARANTEED ANALYSIS**

Crude Protein, %		(min)	20.0
Lysine, %		(min)	1.50
Crude Fat, %		(min)	8.00
Crude Fiber, %		(max)	2.00
Calcium, %	(Ca)	(min)	0.60
Calcium, %	(Ca)	(max)	1.10
Phosphorous, %	(P)	(min)	0.65
Selenium, ppm	(Se)	(min)	0.30
Zinc, ppm	(Zn)	(min)	3,000

#### Medicated for pigs weighing 16-25 lbs.

For the control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to tiamulin and for treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and treatment of bacterial pneumonia caused by Pasteurella multocida susceptible to chlortetracycline.

**Active Drug Ingredient:** Tiamulin (as hydrogen fumarate) 35g/ton Chlortetracycline 400g/ton

### **FEEDING DIRECTIONS**

Offer HSC PRELUDE<sup>m</sup> 309 CTC/DEN to pigs weighing 16 to 25 lbs in bodyweight. Feed continuously as sole ration for 14 days. Feed 0.25 lbs per every 10 of bodyweight to provide 10 mg/lbs bodyweight chlortetracycline.

#### Warning:

Withdraw medicated feed 2 days before slaughter.

#### **Caution:**

Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Semduramicin and Salinomycin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

DO NOT FEED TO SHEEP DUE TO HIGH LEVELS OF COPPER.

#### **Special Considerations:**

Protect from excessive temperatures (temperatures that exceed  $40^{\circ}$  C or  $104^{\circ}$  F). Do not store at warm temperatures up to  $104^{\circ}$  F ( $40^{\circ}$  C) for more than 7 days.