

Offering the Effective Negative Pressure Wound Therapy (NPWT) Patient Care



Easy to Use

- *User-friendly interface*
- *Quickly set pressure from 40-140mmHg*
- *Continuous or Intermittent Modes*
- *Various built-in safety alarms*

Portable

- *Small, lightweight (8.6 oz.) for patient mobility*
- *Disposable canisters that comes in both 110cc and 400cc capacities*
- *Carrying cases available for both 110cc and 400cc sizes*

Re-Chargeable Battery

- *30 hours of continuous battery life*

Dressings

- *Foam, Bandage (Anatomical Fitted) and Foot Dressings. Quick and easy application.*

Casey Lawinski
Customer Care Supervisor
Tel: 215.464.7304
262 Geiger Road Philadelphia, PA 19115
Email: clawinski@pshme.com

Coverage Criteria

Negative Pressure Wound Therapy Pump (E2402) and Supplies (A6550, A7000)

A. Ulcers and Wounds in the Home Setting

The beneficiary has a chronic Stage III or IV pressure ulcer, neuropathic or a chronic (being present for at least 30 days) ulcer of mixed etiology.

or



B. Ulcers and Wounds Encountered in an Inpatient Setting:

An ulcer or wound is encountered in the inpatient setting and, after wound treatments have been tried or considered and ruled out, NPWT is initiated because it is considered in the judgment of the treating physician, the best available treatment option.



**About the medical documentation needed from the patient record,
clinical treatment plan and wound measurements.**